

Dealkylation and Metallation of a Hydrocarbon in On-Surface Synthesis

Wednesday, 26 March 2025 14:14 (3 minutes)

On-Surface Synthesis is established for the creation of extended, one- and two-dimensional macromolecules by homo-coupling directly on a surface [1]. In comparison to classical solution based-chemistry, On-Surface Synthesis has a reduced parameter space (temperature, light) and currently focuses on developing reaction selectivity. Precursor design controls product formation. The build-up of non-stochastic hetero-coupled systems is highly desirable yet not achieved.

Here, we are studying with STM a picene derivative [2] as a precursor, named C12-PicDI, adsorbed on Au (111). Imide moieties, which are reactive to metal ions, are attached to both ends (PicDI). Initially, the reactive end groups are passivated by weakly bond alkyl chains (C12H25). We demonstrate temperature-controlled removal of the alkyls and discuss the system evolution under varying experimental parameters. Alkyl byproducts thermally desorb from and only PicDIs remain on the surface. Selectively removing passivating alkyl groups through controlled heating activates the imide moieties, enabling subsequent metal ion coordination. This work demonstrates a crucial step toward realizing complex, designed heterostructures on surfaces [3].

REFERENCES

- [1] S. Clair and D. G. de Oteyza, Chem. Rev. 119, 4717 (2019).
- [2] Y. Guo, K. Yoshioka, S. Hamao, Y. Kubozono, F. Tani, K. Goto and H. Okamoto, RSC Adv. 10, 31547 (2020).
- [3] G.-Y. Xing, Y.-C. Zhu, D.-Y. Li, and P.-N. Liu, J. Phys. Chem. Lett. 14, 4462 (2023).

Primary authors: AHMED, Aaman Akhtar (National Tsing Hua University); HOFFMANN, Germar (NTHU Dept of Physics)

Co-authors: Prof. OKAMOTO, Hideki (Okayama University); HONG, WEICHENG (NTHU Physics); FENG, Xuan

Presenter: AHMED, Aaman Akhtar (National Tsing Hua University)

Session Classification: Poster Talks

Track Classification: Condensed Matter Experiment